

LINEAR CONNECTIONS AND CURVATURE TENSORS IN THE GEOMETRY OF PARALLELIZABLE MANIFOLDS

Nabil L. Youssef and Amr M. Sid-Ahmed

Department of Mathematics, Faculty of Science,
Cairo University, Giza, Egypt.

nyoussef@frcu.eun.eg and amrs@mailier.eun.eg

Dedicated to the memory of Prof. Dr. A. TAMIM

Abstract. In this paper we discuss linear connections and curvature tensors in the context of geometry of parallelizable manifolds (or absolute parallelism geometry). Different curvature tensors are expressed in a compact form in terms of the torsion tensor of the canonical connection. Using the Bianchi identities, some other identities are derived from the expressions obtained. These identities, in turn, are used to reveal some of the properties satisfied by an intriguing fourth order tensor which we refer to as Wanas tensor.

A further condition on the canonical connection is imposed, assuming it is semi-symmetric. The formulae thus obtained, together with other formulae (Ricci tensors and scalar curvatures of the different connections admitted by the space) are calculated under this additional assumption. Considering a specific form of the semi-symmetric connection causes all nonvanishing curvature tensors to coincide, up to a constant, with the Wanas tensor.

Physical aspects of some of the geometric objects considered are pointed out. ¹

Keywords: Parallelizable manifold, Absolute parallelism geometry, dual connection, semi-symmetric connection, Wanas tensor, Field equations.

AMS Subject Classification. 51P05, 83C05, 53B21.

¹This paper was presented in “The international Conference on Developing and Extending Einstein’s Ideas” held at Cairo, Egypt, December 19-21, 2005.

1. Introduction

After the success of his general theory of gravitation (GR), Einstein searched for a more general theory that would appropriately describe electromagnetic phenomena together with gravity. His search for such a unified theory led him to consider Absolute Parallelism (AP-) geometry [3]. The reason for this is that AP-geometry is wider than the standard Riemannian geometry. According to GR, the ten degrees of freedom (the metric components for $n = 4$) are just sufficient to describe gravitational phenomena alone. AP-geometry, on the other hand, has sixteen degrees of freedom. Riemannian geometry is thus relatively limited compared to AP-geometry which has six extra degrees of freedom. These extra degrees of freedom could be used to describe physical phenomena other than gravity. This idea of increasing the number of degrees of freedom from ten to sixteen is another alternative to the idea of increasing the dimension of the underlying manifold in the so-called Kaluza-Klein theory. Moreover, as opposed to Riemannian geometry which admits only one symmetric affine connection and hence only one curvature tensor, the AP-space admits at least four built-in (natural) affine connections, two of which are non-symmetric. AP-geometry also admits tensors of third order, a number of second order skew and symmetric tensors and a non-vanishing torsion. These extra geometric structures, which have no counterpart in Riemannian geometry, make AP-geometry much richer in its content and hence a potential candidate for geometric unification schemes. A further advantage is that associated to an AP-space there is a Riemannian structure defined in a natural way; thus the AP-space contains within its geometric structure all the mathematical machinery of Riemannian geometry. This facilitates comparison between the results obtained in the context of AP-geometry with the classical GR based on Riemannian geometry. Finally, calculations within the framework of AP-geometry are relatively easier than those used in the context of Riemannian geometry.

In the present paper we investigate the curvature tensors corresponding to the different natural connections defined in an AP-space. The paper is organized in the following manner. In section 2, we give a brief account of the basic elements of AP-geometry; we focus our attention on the fundamental concepts that will be needed in the sequel. In section 3, we consider the curvature tensors of the dual and symmetric connections associated to the canonical connection Γ together with the Riemannian curvature. Simple and compact expressions for such curvature tensors, in terms of the torsion of Γ , are deduced. We then use the Bianchi identities to derive some further interesting identities which simplify the formulae thus obtained. In section 4, we study some of the properties satisfied by an intriguing fourth order tensor, which we call *Wanas tensor* (or simply *W-tensor*) after M.I. Wanas who first defined and used such a tensor [14]. The *W-tensor* is shown to have some properties which are similar to those of the Riemannian curvature tensor. In section 5, we further study the consequences of assuming that the canonical connection is semi-symmetric. All curvature tensors with their associated contractions (Ricci tensors and scalar curvatures) are then derived in this case. We next consider an interesting special case which considerably simplifies the formulae thus obtained and show that all nonvanishing curvature tensors admitted by the space coincide (up to a constant factor) with the *W-tensor*. Physical aspects or physical interpretations of some geometric objects considered are pointed out.

2. A brief account of AP-geometry

In this section, we give a brief account of the geometry of parallelizable manifolds (or absolute parallelism geometry). For more details concerning this geometry, we refer for example to [11], [14], [17] and [2]. Some other works related the subject are [12] and [13].

A parallelizable manifold [2] is an n -dimensional C^∞ manifold M which admits n linearly independent global vector fields λ_i ($i = 1, \dots, n$) on M . Such a space is also known in the literature as absolute parallelism space (AP-space). We will rather use the expressions “AP-space” and “AP-geometry” for their typographical simplicity.

Let λ_i^μ ($\mu = 1, \dots, n$) be the coordinate components of the i -th vector field λ_i . The Einstein summation convention is applied on both Latin (mesh) and Greek (world) indices, where all Latin indices are written as subscripts. In the sequel, we will simply use the symbol λ (without a mesh index) to denote any one of the vector fields λ_i ($i = 1, \dots, n$) and, in most cases, when mesh indices appear they will be in pairs, meaning summation. The covariant components of λ^μ are given via the relations

$$\lambda_i^\mu \lambda_{\nu}^\mu = \delta_{\nu}^\mu, \quad \lambda_i^\mu \lambda_{\mu}^j = \delta_{ij}.$$

The n^3 functions $\Gamma_{\mu\nu}^\alpha$ defined by

$$\Gamma_{\mu\nu}^\alpha := \lambda_i^\alpha \lambda_{\mu,\nu}^i$$

transform as the coefficients of a linear connection under a change of coordinates (where the comma denotes partial differentiation with respect to the coordinate functions). The connection $\Gamma_{\mu\nu}^\alpha$ is clearly non-symmetric and is referred to as the canonical connection of the space. As easily checked, we have

$$\lambda_{\mu|\nu} = 0, \quad \lambda^\mu_{|\nu} = 0,$$

where the stroke denotes covariant differentiation with respect to the canonical connection $\Gamma_{\mu\nu}^\alpha$. The above relation is known in the literature as the condition of absolute parallelism.

Let $\Lambda_{\mu\nu}^\alpha := \Gamma_{\mu\nu}^\alpha - \Gamma_{\nu\mu}^\alpha$ denote the torsion tensor of $\Gamma_{\mu\nu}^\alpha$. It is of particular importance to note that the condition of absolute parallelism together with the commutation formula

$$\lambda^\alpha_{|\mu\nu} - \lambda^\alpha_{|\nu\mu} = \lambda^\epsilon R_{\epsilon\mu\nu}^\alpha + \lambda^\alpha_{|\epsilon} \Lambda_{\nu\mu}^\epsilon$$

forces the curvature tensor $R_{\mu\nu\sigma}^\alpha$ of the canonical connection $\Gamma_{\mu\nu}^\alpha$ to vanish identically. (This is a simple way, which needs no complicated calculations, to prove the vanishing of the curvature tensor $R_{\mu\nu\sigma}^\alpha$). It is for this reason that many authors think that the AP-space is a flat space. This is by no means true. In fact, it is meaningless to speak of curvature without reference to a connection. All we can say is that the AP-space is flat with respect to its canonical connection, or that its canonical connection is flat. However, there are other three natural (built-in) connections which are nonflat. Namely, the dual connection

$$\tilde{\Gamma}_{\mu\nu}^\alpha := \Gamma_{\nu\mu}^\alpha,$$

the symmetric connection

$$\hat{\Gamma}_{\mu\nu}^\alpha := \frac{1}{2}(\Gamma_{\mu\nu}^\alpha + \Gamma_{\nu\mu}^\alpha) = \Gamma_{(\mu\nu)}^\alpha$$

and the Riemannian connection (Christoffel symbols)

$$\overset{\circ}{\Gamma}_{\mu\nu}^{\alpha} := \frac{1}{2} \{g^{\alpha\epsilon} (g_{\epsilon\nu,\mu} + g_{\epsilon\mu,\nu} - g_{\mu\nu,\epsilon})\}$$

associated to the metric structure defined by

$$g_{\mu\nu} := \lambda_{\mu}^{\circ} \lambda_{\nu}^{\circ}$$

with inverse

$$g^{\mu\nu} = \lambda^{\mu}_{\circ} \lambda^{\nu}_{\circ}.$$

It is to be noted that the condition of absolute parallelism implies that the canonical connection $\Gamma_{\mu\nu}^{\alpha}$ is metric :

$$g_{\mu\nu|\sigma} = 0, \quad g^{\mu\nu}{}_{|\sigma} = 0.$$

Consequently, the covariant differentiation defined by the canonical connection commutes with contraction by the metric tensor $g_{\mu\nu}$ and its inverse $g^{\mu\nu}$.

The contortion tensor is defined by

$$\gamma_{\mu\nu}^{\alpha} := \Gamma_{\mu\nu}^{\alpha} - \overset{\circ}{\Gamma}_{\mu\nu}^{\alpha}.$$

Since $\overset{\circ}{\Gamma}_{\mu\nu}^{\alpha}$ is symmetric, it follows that

$$\Lambda_{\mu\nu}^{\alpha} = \gamma_{\mu\nu}^{\alpha} - \gamma_{\nu\mu}^{\alpha}.$$

Moreover, it can be shown that

$$\gamma_{\mu\nu}^{\alpha} = \lambda_{\mu}^{\alpha} \lambda_{\nu}^{\circ}{}_{|\nu},$$

where “ $|$ ” denotes covariant differentiation with respect to the Riemannian connection $\overset{\circ}{\Gamma}_{\mu\nu}^{\alpha}$. Finally, the contortion tensor can be expressed in terms of the torsion tensor in the form [5]:

$$\gamma_{\mu\nu\sigma} = \frac{1}{2} (\Lambda_{\mu\nu\sigma} + \Lambda_{\sigma\nu\mu} + \Lambda_{\nu\sigma\mu}),$$

where $\gamma_{\mu\nu\sigma} = g_{\epsilon\mu} \gamma_{\nu\sigma}^{\epsilon}$ and $\Lambda_{\mu\nu\sigma} = g_{\epsilon\mu} \Lambda_{\nu\sigma}^{\epsilon}$. It is to be noted that $\Lambda_{\mu\nu\sigma}$ is skew-symmetric in the last pair of indices, whereas $\gamma_{\mu\nu\sigma}$ is skew-symmetric in the first pair of indices. It is to be noted also that the contortion tensor vanishes if and only if the torsion tensor vanishes.

We have four types of covariant derivatives corresponding to the four connections mentioned above, namely

$$A^{\mu}{}_{|\nu} = A^{\mu}{}_{,\nu} + \Gamma_{\epsilon\nu}^{\mu} A^{\epsilon},$$

$$A^{\mu}{}_{\tilde{|\nu}} = A^{\mu}{}_{,\nu} + \tilde{\Gamma}_{\epsilon\nu}^{\mu} A^{\epsilon},$$

$$A^{\mu}{}_{\hat{|\nu}} = A^{\mu}{}_{,\nu} + \hat{\Gamma}_{\epsilon\nu}^{\mu} A^{\epsilon},$$

$$A^{\mu}{}_{\circ|\nu} = A^{\mu}{}_{,\nu} + \overset{\circ}{\Gamma}_{\epsilon\nu}^{\mu} A^{\epsilon},$$

where A^{μ} is an arbitrary contravariant vector.

In conclusion, the AP-space has four curvature tensors $R_{\mu\nu\sigma}^{\alpha}$, $\tilde{R}_{\mu\nu\sigma}^{\alpha}$, $\hat{R}_{\mu\nu\sigma}^{\alpha}$ and $\overset{\circ}{R}_{\mu\nu\sigma}^{\alpha}$ corresponding to the four connections $\Gamma_{\mu\nu}^{\alpha}$, $\tilde{\Gamma}_{\mu\nu}^{\alpha}$, $\hat{\Gamma}_{\mu\nu}^{\alpha}$ and $\overset{\circ}{\Gamma}_{\mu\nu}^{\alpha}$ respectively. As already mentioned, only one of these curvature tensors vanishes identically (the curvature $R_{\mu\nu\sigma}^{\alpha}$ of the canonical connection $\Gamma_{\mu\nu}^{\alpha}$). The other three do not vanish in general. The vanishing of $R_{\mu\nu\sigma}^{\alpha}$ enables us to express the other three curvature tensors in terms of the torsion tensor $\Lambda_{\mu\nu}^{\alpha}$ as it will be revealed in the next section.

Summary of the geometry of the AP-space

Connection	Coefficients	Covariant derivative	Torsion	Curvature	Metricity
Canonical	$\Gamma_{\mu\nu}^\alpha$	$ $	$\Lambda_{\mu\nu}^\alpha$	0	metric
Dual	$\tilde{\Gamma}_{\mu\nu}^\alpha$	$\sim $	$-\Lambda_{\mu\nu}^\alpha$	$\tilde{R}_{\mu\nu\sigma}^\alpha$	non-metric
Symmetric	$\hat{\Gamma}_{\mu\nu}^\alpha$	$\hat{ }$	0	$\hat{R}_{\mu\nu\sigma}^\alpha$	non-metric
Riemannian	$\overset{\circ}{\Gamma}_{\mu\nu}^\alpha$	$\overset{\circ}{ }$	0	$\overset{\circ}{R}_{\mu\nu\sigma}^\alpha$	metric

3. Curvature tensors and the Bianchi identities in AP-geometry

Let (M, λ) be an AP-space of dimension n , where λ denotes any one of the n linearly independent vector fields λ_i ($i = 1, \dots, n$) defining the AP-structure on M . Let $\Gamma_{\mu\nu}^\alpha$ be the canonical connection on M defined by $\Gamma_{\mu\nu}^\alpha = \lambda_i^\alpha \lambda_{i\mu\nu}$. Let $\tilde{\Gamma}_{\mu\nu}^\alpha$, $\hat{\Gamma}_{\mu\nu}^\alpha$ and $\overset{\circ}{\Gamma}_{\mu\nu}^\alpha$ be respectively the dual connection associated to $\Gamma_{\mu\nu}^\alpha$, the symmetric connection associated to $\Gamma_{\mu\nu}^\alpha$ and the Riemannian connection defined by the metric tensor $g_{\mu\nu} = \lambda_{i\mu} \lambda_{i\nu}$. As in the previous section, covariant differentiation with respect to $\Gamma_{\mu\nu}^\alpha$, $\tilde{\Gamma}_{\mu\nu}^\alpha$, $\hat{\Gamma}_{\mu\nu}^\alpha$ and $\overset{\circ}{\Gamma}_{\mu\nu}^\alpha$ will be denoted by $|$, $\sim|$, $\hat{|}$ and $\overset{\circ}{|}$ respectively.

Theorem 3.1. *The curvature tensors $\tilde{R}_{\mu\nu\sigma}^\alpha$, $\hat{R}_{\mu\nu\sigma}^\alpha$ and $\overset{\circ}{R}_{\mu\nu\sigma}^\alpha$ of the connections $\tilde{\Gamma}_{\mu\nu}^\alpha$, $\hat{\Gamma}_{\mu\nu}^\alpha$ and $\overset{\circ}{\Gamma}_{\mu\nu}^\alpha$ are expressed in terms of the torsion tensor $\Lambda_{\mu\nu}^\alpha$ of the canonical connection $\Gamma_{\mu\nu}^\alpha$ as follows:*

- (a) $\tilde{R}_{\mu\nu\sigma}^\alpha = \Lambda_{\sigma\nu|\mu}^\alpha$.
- (b) $\hat{R}_{\mu\nu\sigma}^\alpha = \frac{1}{2}(\Lambda_{\mu\nu|\sigma}^\alpha - \Lambda_{\mu\sigma|\nu}^\alpha) + \frac{1}{4}(\Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha - \Lambda_{\mu\sigma}^\epsilon \Lambda_{\nu\epsilon}^\alpha) + \frac{1}{2}(\Lambda_{\sigma\nu}^\epsilon \Lambda_{\epsilon\mu}^\alpha)$.
- (c) $\overset{\circ}{R}_{\mu\nu\sigma}^\alpha = (\gamma_{\mu\nu|\sigma}^\alpha - \gamma_{\mu\sigma|\nu}^\alpha) + (\gamma_{\mu\sigma}^\epsilon \gamma_{\epsilon\nu}^\alpha - \gamma_{\mu\nu}^\epsilon \gamma_{\epsilon\sigma}^\alpha) + \gamma_{\mu\epsilon}^\alpha \Lambda_{\nu\sigma}^\epsilon$.

Proof. We start by proving the first relation:

$$\begin{aligned}
\tilde{R}_{\mu\nu\sigma}^\alpha &= \tilde{\Gamma}_{\mu\sigma,\nu}^\alpha - \tilde{\Gamma}_{\mu\nu,\sigma}^\alpha + \tilde{\Gamma}_{\mu\sigma}^\epsilon \tilde{\Gamma}_{\epsilon\nu}^\alpha - \tilde{\Gamma}_{\mu\nu}^\epsilon \tilde{\Gamma}_{\epsilon\sigma}^\alpha \\
&= \Gamma_{\sigma\mu,\nu}^\alpha - \Gamma_{\nu\mu,\sigma}^\alpha + \Gamma_{\sigma\mu}^\epsilon \Gamma_{\nu\epsilon}^\alpha - \Gamma_{\nu\mu}^\epsilon \Gamma_{\sigma\epsilon}^\alpha \\
&= \{\Gamma_{\sigma\mu,\nu}^\alpha + \Gamma_{\sigma\mu}^\epsilon (\Lambda_{\nu\epsilon}^\alpha + \Gamma_{\nu\epsilon}^\alpha)\} - \{\Gamma_{\nu\mu,\sigma}^\alpha + \Gamma_{\nu\mu}^\epsilon (\Lambda_{\sigma\epsilon}^\alpha + \Gamma_{\sigma\epsilon}^\alpha)\} \\
&= (\Gamma_{\sigma\mu,\nu}^\alpha + \Gamma_{\sigma\mu}^\epsilon \Gamma_{\epsilon\nu}^\alpha) - (\Gamma_{\nu\mu,\sigma}^\alpha + \Gamma_{\nu\mu}^\epsilon \Gamma_{\epsilon\sigma}^\alpha) - (\Gamma_{\sigma\mu}^\epsilon \Lambda_{\epsilon\nu}^\alpha + \Gamma_{\nu\mu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha) \\
&= (R_{\sigma\nu\mu}^\alpha + \Gamma_{\sigma\nu,\mu}^\epsilon + \Gamma_{\sigma\nu}^\epsilon \Gamma_{\epsilon\mu}^\alpha) + (R_{\nu\mu\sigma}^\alpha - \Gamma_{\nu\sigma,\mu}^\epsilon - \Gamma_{\nu\sigma}^\epsilon \Gamma_{\epsilon\mu}^\alpha) - (\Gamma_{\sigma\mu}^\epsilon \Lambda_{\epsilon\nu}^\alpha + \Gamma_{\nu\mu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha).
\end{aligned}$$

Taking into account the fact that $R_{\mu\nu\sigma}^\alpha = 0$, we get

$$\tilde{R}_{\mu\nu\sigma}^\alpha = \Lambda_{\sigma\nu,\mu}^\alpha + \Gamma_{\epsilon\mu}^\alpha \Lambda_{\sigma\nu}^\epsilon - \Gamma_{\sigma\mu}^\epsilon \Lambda_{\epsilon\nu}^\alpha - \Gamma_{\nu\mu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha = \Lambda_{\sigma\nu|\mu}^\alpha.$$

We then prove the second relation: We have, by definition,

$$\hat{R}_{\mu\nu\sigma}^\alpha = \hat{\Gamma}_{\mu\sigma,\nu}^\alpha - \hat{\Gamma}_{\mu\nu,\sigma}^\alpha + \hat{\Gamma}_{\mu\sigma}^\epsilon \hat{\Gamma}_{\epsilon\nu}^\alpha - \hat{\Gamma}_{\mu\nu}^\epsilon \hat{\Gamma}_{\epsilon\sigma}^\alpha.$$

Now,

$$\begin{aligned}
\hat{\Gamma}_{\mu\sigma}^\epsilon \hat{\Gamma}_{\epsilon\nu}^\alpha &= \frac{1}{4}(\Lambda_{\sigma\mu}^\epsilon + 2\Gamma_{\mu\sigma}^\epsilon)(\Lambda_{\nu\epsilon}^\alpha + 2\Gamma_{\epsilon\nu}^\alpha) \\
&= -\frac{1}{4}\Lambda_{\mu\sigma}^\epsilon \Lambda_{\nu\epsilon}^\alpha - \frac{1}{2}\Lambda_{\mu\sigma}^\epsilon \Gamma_{\epsilon\nu}^\alpha - \frac{1}{2}\Lambda_{\epsilon\nu}^\alpha \Gamma_{\mu\sigma}^\epsilon + \Gamma_{\mu\sigma}^\epsilon \Gamma_{\epsilon\nu}^\alpha.
\end{aligned}$$

Similarly,

$$\hat{\Gamma}_{\mu\nu}^\epsilon \hat{\Gamma}_{\epsilon\sigma}^\alpha = -\frac{1}{4}\Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha - \frac{1}{2}\Lambda_{\mu\nu}^\epsilon \Gamma_{\epsilon\sigma}^\alpha - \frac{1}{2}\Lambda_{\epsilon\sigma}^\alpha \Gamma_{\mu\nu}^\epsilon + \Gamma_{\mu\nu}^\epsilon \Gamma_{\epsilon\sigma}^\alpha.$$

Moreover,

$$\hat{\Gamma}_{\mu\sigma,\nu}^\alpha = -\frac{1}{2}\Lambda_{\mu\sigma,\nu}^\alpha + \Gamma_{\mu\sigma,\nu}^\alpha \quad \text{and} \quad \hat{\Gamma}_{\mu\nu,\sigma}^\alpha = -\frac{1}{2}\Lambda_{\mu\nu,\sigma}^\alpha + \Gamma_{\mu\nu,\sigma}^\alpha.$$

Hence, noting that $R_{\mu\nu\sigma}^\alpha = 0$, we get

$$\begin{aligned}
\hat{R}_{\mu\nu\sigma}^\alpha &= \frac{1}{4}(\Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha - \Lambda_{\mu\sigma}^\epsilon \Lambda_{\nu\epsilon}^\alpha) + \frac{1}{2}\{(\Lambda_{\mu\nu,\sigma}^\alpha + \Lambda_{\mu\nu}^\epsilon \Gamma_{\epsilon\sigma}^\alpha - \Lambda_{\epsilon\nu}^\alpha \Gamma_{\mu\sigma}^\epsilon) \\
&\quad - (\Lambda_{\mu\sigma,\nu}^\alpha + \Lambda_{\mu\sigma}^\epsilon \Gamma_{\epsilon\nu}^\alpha - \Lambda_{\epsilon\sigma}^\alpha \Gamma_{\mu\nu}^\epsilon)\} \\
&= \frac{1}{2}(\Lambda_{\mu\nu|\sigma}^\alpha - \Lambda_{\mu\sigma|\nu}^\alpha) + \frac{1}{4}(\Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha - \Lambda_{\mu\sigma}^\epsilon \Lambda_{\nu\epsilon}^\alpha) + \frac{1}{2}(\Lambda_{\sigma\nu}^\epsilon \Lambda_{\epsilon\mu}^\alpha).
\end{aligned}$$

The proof of relation (c) is carried out in the same manner and we omit it. $\square$

Remark 3.2. *The first and second formulae (resp. The third formula) of the above theorem remain (resp. remains) valid in the more general context in which $\Gamma_{\mu\nu}^\alpha$ is any given non-symmetric linear connection on a manifold M (resp. a Riemannian manifold (M, g)) with vanishing curvature.*

It is clear that the torsion tensor plays the key role in all identities obtained above. The vanishing of the torsion tensor forces the three connections $\Gamma_{\mu\nu}^\alpha$, $\tilde{\Gamma}_{\mu\nu}^\alpha$ and $\hat{\Gamma}_{\mu\nu}^\alpha$ to coincide with the Riemannian connection $\overset{\circ}{\Gamma}_{\mu\nu}^\alpha$ and the AP-space in this case becomes trivially a flat Riemannian space. Consequently, the non-vanishing of any of the three curvature tensors suffices for the non-vanishing of the torsion tensor.

We now derive some relations that will prove useful later on.

Proposition 3.3. *The following relations hold:*

- (a) $\Lambda_{\mu\nu|\sigma}^\alpha - \Lambda_{\mu\nu\tilde{|\sigma}}^\alpha = \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu}^\epsilon \Lambda_{\epsilon\sigma}^\alpha.$
- (b) $\Lambda_{\mu\nu|\sigma}^\alpha - \Lambda_{\mu\nu\hat{|\sigma}}^\alpha = \frac{1}{2}(\mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu}^\epsilon \Lambda_{\epsilon\sigma}^\alpha).$
- (c) $\Lambda_{\mu\nu|\sigma}^\alpha - \Lambda_{\mu\nu\circ\sigma}^\alpha = \Lambda_{\mu\nu}^\epsilon \gamma_{\epsilon\sigma}^\alpha + \Lambda_{\nu\epsilon}^\alpha \gamma_{\mu\sigma}^\epsilon + \Lambda_{\epsilon\mu}^\alpha \gamma_{\nu\sigma}^\epsilon.$

where the notation $\mathfrak{S}_{\mu,\nu,\sigma}$ denotes a cyclic permutation of the indices μ, ν, σ and summation.

Proof. The three relations follow from the definition of the covariant derivative with respect to the appropriate connection. $\square$

Let M be a differentiable manifold equipped with a linear connection with torsion T and curvature R . Then the Bianchi identities are given locally by [8]:

$$\mathfrak{S}_{\mu,\nu,\sigma} R_{\mu\nu\sigma}^\alpha = \mathfrak{S}_{\mu,\nu,\sigma} (T_{\mu\nu;\sigma}^\alpha + T_{\mu\nu}^\epsilon T_{\epsilon\sigma}^\alpha), \text{ (first Bianchi identity)}$$

$$\mathfrak{S}_{\mu,\nu,\sigma} (R_{\beta\mu\nu;\sigma}^\alpha + R_{\beta\mu\epsilon}^\alpha T_{\nu\sigma}^\epsilon) = 0. \quad \text{(second Bianchi identity)}$$

where “;” denotes covariant differentiation with respect to the given connection.

In what follows, we derive some identities using the first and second Bianchi identities. Some of the derived identities will be used in simplifying some of the formulae thus obtained.

Theorem 3.4. *The first Bianchi identity for the connections $\Gamma_{\mu\nu}^\alpha$, $\tilde{\Gamma}_{\mu\nu}^\alpha$, $\hat{\Gamma}_{\mu\nu}^\alpha$ and $\mathring{\Gamma}_{\mu\nu}^\alpha$ reads:*

- (a) $\mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\mu\nu|\sigma}^\alpha + \Lambda_{\mu\nu}^\epsilon \Lambda_{\epsilon\sigma}^\alpha) = 0.$
- (b) $\mathfrak{S}_{\mu,\nu,\sigma} \tilde{R}_{\mu\nu\sigma}^\alpha = \mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\nu\mu\tilde{|\sigma}}^\alpha + \Lambda_{\mu\nu}^\epsilon \Lambda_{\epsilon\sigma}^\alpha).$
- (c) $\mathfrak{S}_{\mu,\nu,\sigma} \hat{R}_{\mu\nu\sigma}^\alpha = 0.$
- (d) $\mathfrak{S}_{\mu,\nu,\sigma} \mathring{R}_{\mu\nu\sigma}^\alpha = 0.$

The second Bianchi identity for the connections $\tilde{\Gamma}_{\mu\nu}^\alpha$, $\hat{\Gamma}_{\mu\nu}^\alpha$ and $\mathring{\Gamma}_{\mu\nu}^\alpha$ reads:

- (e) $\mathfrak{S}_{\mu,\nu,\sigma} \tilde{R}_{\beta\mu\nu\tilde{|\sigma}}^\alpha = \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\sigma\nu}^\epsilon \Lambda_{\epsilon\mu|\beta}^\alpha.$
- (f) $\mathfrak{S}_{\mu,\nu,\sigma} \hat{R}_{\beta\mu\nu\hat{|\sigma}}^\alpha = 0.$
- (g) $\mathfrak{S}_{\mu,\nu,\sigma} \mathring{R}_{\beta\mu\nu\mathring{|\sigma}}^\alpha = 0.$

Proof. Identities (a) and (b) follow respectively from the fact that $R_{\mu\nu\sigma}^\alpha$ vanishes identically and that $\tilde{\Lambda}_{\mu\nu}^\alpha = \Lambda_{\nu\mu}^\alpha$. Identity (e) results from Theorem 3.1 (a) together with the fact that $\tilde{\Lambda}_{\mu\nu}^\alpha = \Lambda_{\nu\mu}^\alpha$. The remaining identities are trivial because of the symmetry of the connections $\hat{\Gamma}_{\mu\nu}^\alpha$ and $\mathring{\Gamma}_{\mu\nu}^\alpha$. $\square$

Proposition 3.5. *The following identities hold:*

- (a) $\mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu|\sigma}^\alpha = 0.$
- (b) $\mathfrak{S}_{\mu,\nu,\sigma} \tilde{R}_{\mu\nu\sigma}^\alpha = \mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\mu\nu}^\epsilon \Lambda_{\epsilon\sigma}^\alpha).$

Proof. Taking into account Theorem 3.4 (b) together with Theorem 3.1 (a) we get

$$\mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\nu\sigma|\mu}^\alpha + \Lambda_{\mu\nu}^\epsilon \Lambda_{\epsilon\sigma}^\alpha) = \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu|\sigma}^\alpha.$$

By Theorem 3.4 (a), the left hand side of the the above equation vanishes and the first identity follows. The second identity is a direct consequence of theorem 3.4 (b), taking into consideration identity (a) above. $\square$

The next result is crucial in simplifying some of the identities obtained so far and in proving other interesting results.

Theorem 3.6. *The torsion tensor satisfies the identity*

$$\mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu}^\epsilon \Lambda_{\epsilon\sigma}^\alpha = 0.$$

Proof. Applying the first Bianchi identity to $\hat{R}_{\mu\nu\sigma}^\alpha$ as expressed in Theorem 3.1 (b), we get

$$\frac{1}{2} \mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\mu\nu|\sigma}^\alpha - \Lambda_{\mu\sigma|\nu}^\alpha) + \frac{1}{4} \mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha - \Lambda_{\mu\sigma}^\epsilon \Lambda_{\nu\epsilon}^\alpha) + \frac{1}{2} \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\nu\sigma}^\epsilon \Lambda_{\mu\epsilon}^\alpha = 0. \quad (*)$$

Considering each of the above three terms separately, and taking into account Theorem 3.4 (a), we obtain respectively

$$\begin{aligned} \frac{1}{2} \mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\mu\nu|\sigma}^\alpha - \Lambda_{\mu\sigma|\nu}^\alpha) &= \frac{1}{2} \mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\mu\nu|\sigma}^\alpha + \Lambda_{\sigma\mu|\nu}^\alpha) = \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu|\sigma}^\alpha = \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha, \\ \frac{1}{4} \mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha - \Lambda_{\mu\sigma}^\epsilon \Lambda_{\nu\epsilon}^\alpha) &= \frac{1}{4} \mathfrak{S}_{\mu,\nu,\sigma} (\Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha + \Lambda_{\sigma\mu}^\epsilon \Lambda_{\nu\epsilon}^\alpha) = \frac{1}{2} \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha \end{aligned}$$

and

$$\frac{1}{2} \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\nu\sigma}^\epsilon \Lambda_{\mu\epsilon}^\alpha = \frac{1}{2} \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu}^\epsilon \Lambda_{\sigma\epsilon}^\alpha.$$

The required identity results by substituting the above three equations into (*). $\square$

Corollary 3.7. *The following identities hold:*

- (a) $\Lambda_{\mu\nu|\sigma}^\alpha = \Lambda_{\mu\nu|\sigma}^\alpha = \Lambda_{\mu\nu|\sigma}^\alpha.$
- (b) $\mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\mu\nu|\sigma}^\alpha = 0.$
- (c) $\mathfrak{S}_{\mu,\nu,\sigma} \tilde{R}_{\mu\nu\sigma}^\alpha = 0.$
- (d) $\mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\nu\mu|\beta\sigma}^\alpha = \mathfrak{S}_{\mu,\nu,\sigma} \Lambda_{\sigma\nu}^\epsilon \Lambda_{\epsilon\mu|\beta}^\alpha \quad (\Lambda_{\nu\mu|\beta\sigma}^\alpha := \Lambda_{\nu\mu|\beta|\sigma}^\alpha).$
- (e) $\mathfrak{S}_{\mu,\nu,\sigma} (\gamma_{\mu\nu|\sigma}^\alpha - \gamma_{\nu\mu|\sigma}^\alpha) = 0.$

Proof. Taking into account Theorem 3.6, relation (a) follows from Proposition 3.3 (a) and (b), whereas identities (b) and (c) follow from Theorem 3.4 (a) and Proposition 3.5 (b) respectively. Identity (d) follows from Theorem 3.4 (e) taking into account Theorem 3.1 (a) together with relation (a) above. Finally, the last identity follows from identity (b) above together with the relation $\Lambda_{\mu\nu}^\alpha = \gamma_{\mu\nu}^\alpha - \gamma_{\nu\mu}^\alpha$. $\square$

In view of Theorem 3.6, the curvature tensor $\hat{R}_{\mu\nu\sigma}^\alpha$ can be further simplified as revealed in

Proposition 3.8. *The curvature tensor $\hat{R}_{\mu\nu\sigma}^\alpha$ can be expressed in the form:*

$$\hat{R}_{\mu\nu\sigma}^\alpha = \frac{1}{2}\Lambda_{\sigma\nu|\mu}^\alpha + \frac{1}{4}\Lambda_{\sigma\nu}^\epsilon\Lambda_{\epsilon\mu}^\alpha.$$

Proof. The curvature tensor $\hat{R}_{\mu\nu\sigma}^\alpha$ has the form (Theorem 3.1 (b)):

$$\hat{R}_{\mu\nu\sigma}^\alpha = \frac{1}{2}(\Lambda_{\mu\nu|\sigma}^\alpha - \Lambda_{\mu\sigma|\nu}^\alpha) + \frac{1}{4}(\Lambda_{\mu\nu}^\epsilon\Lambda_{\sigma\epsilon}^\alpha - \Lambda_{\mu\sigma}^\epsilon\Lambda_{\nu\epsilon}^\alpha) + \frac{1}{2}(\Lambda_{\sigma\nu}^\epsilon\Lambda_{\epsilon\mu}^\alpha).$$

Taking into account identity (b) in Corollary 3.7 and Theorem 3.6, we get

$$\begin{aligned}\hat{R}_{\mu\nu\sigma}^\alpha &= \frac{1}{2}\{(\mathfrak{S}_{\mu,\nu,\sigma}\Lambda_{\mu\nu|\sigma}^\alpha) - \Lambda_{\nu\sigma|\mu}^\alpha\} + \frac{1}{2}\Lambda_{\nu\sigma}^\epsilon\Lambda_{\epsilon\mu}^\alpha \\ &\quad + \frac{1}{4}\{(\mathfrak{S}_{\mu,\nu,\epsilon}\Lambda_{\mu\nu}^\epsilon\Lambda_{\sigma\epsilon}^\alpha) - \Lambda_{\nu\sigma}^\epsilon\Lambda_{\mu\epsilon}^\alpha\} \\ &= \frac{1}{2}\Lambda_{\sigma\nu|\mu}^\alpha + \frac{1}{4}\Lambda_{\sigma\nu}^\epsilon\Lambda_{\epsilon\mu}^\alpha. \quad \square\end{aligned}$$

The formula obtained in Theorem 3.1 for the curvature tensor $\tilde{R}_{\mu\nu\sigma}^\alpha$ is strikingly compact and elegant. The formula obtained for $\hat{R}_{\mu\nu\sigma}^\alpha$ is however less elegant but still relatively compact. These two formulae, together with the first Bianchi identity, enabled us to derive, in a simple way, the crucial identities $\mathfrak{S}_{\mu,\nu,\sigma}\Lambda_{\mu\nu|\sigma}^\alpha = 0$ and $\mathfrak{S}_{\mu,\nu,\sigma}\Lambda_{\mu\nu}^\epsilon\Lambda_{\sigma\epsilon}^\alpha = 0$ which, in turn, simplified the formula obtained for $\hat{R}_{\mu\nu\sigma}^\alpha$ (which is now more elegant). The last two identities will play an essential role in the rest of the paper. It should be noted however that a direct proof of these identities is far from clear.

Remark 3.9. *All results and identities concerning the connections $\Gamma_{\mu\nu}^\alpha$, $\tilde{\Gamma}_{\mu\nu}^\alpha$ and $\hat{\Gamma}_{\mu\nu}^\alpha$ (resp. the connection $\mathring{\Gamma}_{\mu\nu}^\alpha$) remain valid in the more general context in which $\Gamma_{\mu\nu}^\alpha$ is a non-symmetric linear connection on a manifold M (resp. a Riemannian manifold (M, g)) with vanishing curvature.*

4. The Wanas Tensor (W-tensor)

Let (M, λ) be an AP-space of dimension n , where λ denotes any one of the n linearly independent vector fields defining the AP-structure on the manifold M . Let $\tilde{R}_{\mu\nu\sigma}^\alpha$ and $\tilde{\Lambda}_{\mu\nu}^\alpha$ be the curvature and the torsion tensors of the dual connection $\tilde{\Gamma}_{\mu\nu}^\alpha$.

Definition 4.1. The tensor field $W_{\mu\nu\sigma}^\alpha$ of type $(1,3)$ on M defined by the formula

$$\lambda_\epsilon W_{\mu\nu\sigma}^\epsilon := \lambda_{\mu|\nu\sigma} - \lambda_{\mu|\sigma\nu}$$

will be called the Wanas tensor, or simply the W -tensor, of the AP -space (M, λ) .

The W -tensor $W_{\mu\nu\sigma}^\alpha$ has been first defined by M. I Wanas [14] in 1975 and has been used by F. Mikhail and M. Wanas [9] to construct a pure geometric theory unifying gravity and electromagnetism. A double contraction of the tensor $g^{\mu\gamma} W_{\mu\nu\sigma}^\alpha$ gives a scalar that has been used to write the Lagrangian density for fields and matter. The symmetric part of the field equations obtained contains a second order tensor representing the material distribution. This tensor is a pure geometric object and not a phenomenological one. The skew part of the field equations gives rise to Maxwell-like equations. In this theory the metric tensor plays the role of the gravitational potential while the the vector c_μ (the contracted torsion) plays the role of the electromagnetic potential. The linearized form [10] of the theory supports these identifications.

The next result gives quite a simple expression for such a tensor.

Theorem 4.2. Let (M, λ) be an AP -space. Then the W -tensor can be expressed in the form

$$W_{\mu\nu\sigma}^\alpha = \Lambda_{\sigma\nu|\mu}^\alpha - \Lambda_{\sigma\nu}^\epsilon \Lambda_{\epsilon\mu}^\alpha,$$

where $\Lambda_{\mu\nu}^\alpha$ is the torsion tensor of the canonical connection $\Gamma_{\mu\nu}^\alpha$.

Proof. Consider the commutation formula with respect to the connection $\tilde{\Gamma}_{\mu\nu}^\alpha$:

$$\lambda_{\mu|\nu\sigma} - \lambda_{\mu|\sigma\nu} = \lambda_\epsilon \tilde{R}_{\mu\nu\sigma}^\epsilon + \lambda_{\mu|\epsilon} \tilde{\Lambda}_{\nu\sigma}^\epsilon.$$

Multiplying both sides by λ^α , using the definition of the W -tensor together with the definition of $\Gamma_{\mu\nu}^\alpha$ and taking into account Theorem 3.1 (a), we get

$$\begin{aligned} W_{\mu\nu\sigma}^\alpha &= \Lambda_{\sigma\nu|\mu}^\alpha + \lambda_i^\alpha (\lambda_{\mu,\epsilon} - \lambda_{\epsilon\mu}) \Lambda_{\sigma\nu}^\epsilon \\ &= \Lambda_{\sigma\nu|\mu}^\alpha + (\Gamma_{\mu\epsilon}^\alpha - \Gamma_{\epsilon\mu}^\alpha) \Lambda_{\sigma\nu}^\epsilon \\ &= \Lambda_{\sigma\nu|\mu}^\alpha - \Lambda_{\sigma\nu}^\epsilon \Lambda_{\epsilon\mu}^\alpha. \quad \square \end{aligned}$$

Remark 4.3. The W -tensor can be also defined contravariantly as follows:

$$\lambda^\mu W_{\mu\nu\sigma}^\alpha = \lambda^\alpha \tilde{\lambda}_{\nu\sigma} - \lambda^\alpha \tilde{\lambda}_{\sigma\nu} = \lambda^\epsilon \tilde{R}_{\epsilon\sigma\nu}^\alpha + \lambda^\alpha \tilde{\lambda}_{\epsilon} \tilde{\Lambda}_{\nu\sigma}^\epsilon.$$

This definition gives the same formula for the W -tensor as in Theorem 4.2.

Proposition 4.4. The Wanas tensor has the following properties:

a) $W_{\mu\nu\sigma}^\alpha$ is skew symmetric in the last pair of indices.

(b) $W_{\mu\nu\sigma|\beta}^\alpha - W_{\mu\nu\sigma\tilde{\beta}}^\alpha = \Lambda_{\sigma\nu|\mu\beta}^\alpha - \Lambda_{\sigma\nu\tilde{\mu}\beta}^\alpha$.

Proof. The first property is trivial. The second property holds as a result of Theorem 4.2 together with corollary 3.7 (a). $\square$

Theorem 4.5. *The W-tensor satisfies the following identity:*

$$\mathfrak{S}_{\mu,\nu,\sigma} W_{\mu\nu\sigma}^\alpha = 0.$$

Proof. Follows directly from Theorem 4.2, taking into account Theorem 3.6 together with Corollary 3.7 (b). $\square$

The identity satisfied by the W-tensor in Theorem 4.5 is the same as the first Bianchi identity of the Riemannian curvature tensor. The identity corresponding to the second Bianchi identity is given by:

Theorem 4.6. *The W-tensor satisfies the following identity:*

$$\mathfrak{S}_{\nu,\sigma,\beta} W_{\mu\nu\sigma|\beta}^\alpha = \mathfrak{S}_{\nu,\sigma,\beta} \{ \Lambda_{\nu\sigma}^\epsilon (\Lambda_{\epsilon\beta|\mu}^\alpha + \Lambda_{\epsilon\mu|\beta}^\alpha) \}.$$

Proof. Taking into account the second Bianchi identity, Theorem 4.2, Theorem 3.1 (a) and Corollary 3.8 (a) and (b), we get

$$\begin{aligned} \mathfrak{S}_{\nu,\sigma,\beta} W_{\mu\nu\sigma|\beta}^\alpha &= \mathfrak{S}_{\nu,\sigma,\beta} \{ \tilde{R}_{\mu\nu\sigma|\beta}^\alpha - (\Lambda_{\sigma\nu}^\epsilon \Lambda_{\epsilon\mu}^\alpha)_{|\beta} \} \\ &= \mathfrak{S}_{\nu,\sigma,\beta} (\tilde{R}_{\mu\nu\epsilon}^\alpha \tilde{\Lambda}_{\beta\sigma}^\epsilon - \Lambda_{\sigma\nu}^\epsilon \Lambda_{\epsilon\mu|\beta}^\alpha) \\ &= \mathfrak{S}_{\nu,\sigma,\beta} (\Lambda_{\epsilon\nu|\mu}^\alpha \Lambda_{\sigma\beta}^\epsilon - \Lambda_{\epsilon\mu|\beta}^\alpha \Lambda_{\sigma\nu}^\epsilon) \\ &= \mathfrak{S}_{\nu,\sigma,\beta} \{ \Lambda_{\nu\sigma}^\epsilon (\Lambda_{\epsilon\beta|\mu}^\alpha + \Lambda_{\epsilon\mu|\beta}^\alpha) \}. \quad \square \end{aligned}$$

We conclude this section by some comments :

- The W-tensor is defined in terms of the dual connection $\tilde{\Gamma}_{\mu\nu}^\alpha$. The same definition using the three other connections gives nothing new. In fact, the commutation formula for the connection $\Gamma_{\nu\mu}^\alpha$ is trivial (since $\lambda^\alpha_{|\mu} = 0$), whereas the commutation formulae for $\hat{\Gamma}_{\mu\nu}^\alpha$ and $\overset{\circ}{\Gamma}_{\mu\nu}^\alpha$ give rise to $\hat{R}_{\nu\mu\sigma}^\alpha$ and $\overset{\circ}{R}_{\mu\nu\sigma}^\alpha$ respectively (since the torsion tensor vanishes in the latter two cases).
- The vanishing of the torsion tensor implies the vanishing of the W-tensor which is equivalent to the commutativity of successive covariant differentiation (of the vector fields λ); a striking property which does not exist in Riemannian geometry (or even in other geometries, in general).
- The W-tensor has some properties common with the Riemannian curvature tensor (for example, Proposition 4.4 (a) and Theorem 4.5). Nevertheless, there are significant deviations from the Riemannian curvature tensor (for example, Theorem 4.6).
- As stated above, Mikhail and Wanas [9], in their theory, have attributed the gravitational potential to the metric tensor and the electromagnetic potential to the contracted torsion. The use of the W-tensor has aided to construct such a theory **via one single geometric entity**, which Einstein was seeking for [4]. Different applications (for example, [15], [16]) of Mikhail-Wanas theory support these statements. In conclusion, the expression of the W-tensor comprises, in addition to the curvature tensor $\tilde{R}_{\mu\nu\sigma}^\alpha$, a term containing a torsion contribution. Thus, the W-tensor expresses *geometrically* the interaction between curvature and torsion. On the other hand, as gravity is described in terms of curvature and electromagnetism is described in terms of torsion, the W-tensor expresses *physically* the interaction between gravity and electromagnetism.

5. The semi-symmetric case

In this section, we investigate an AP-space whose canonical connection has a special simple form.

Definition 5.1. *Let M be a differentiable manifold. A semi-symmetric connection on M is a linear connection on M whose torsion tensor T is given by [18]*

$$T_{\mu\nu}^\alpha = \delta_\mu^\alpha w_\nu - \delta_\nu^\alpha w_\mu,$$

for some scalar 1-form w_μ .

Semi-symmetric connections have been studied by many authors (cf. for example [1], [6], [19], [20]). In what follows, we consider an n -dimensional AP-space (M, λ) with the additional assumption that the canonical connection $\Gamma_{\mu\nu}^\alpha$ is semi-symmetric. Then, by Definition 5.1, we have

$$\Lambda_{\mu\nu}^\alpha = \delta_\mu^\alpha w_\nu - \delta_\nu^\alpha w_\mu.$$

Moreover, it can be shown that [7]

$$\Gamma_{\mu\nu}^\alpha = \overset{\circ}{\Gamma}_{\mu\nu}^\alpha + \delta_\mu^\alpha w_\nu - g_{\mu\nu} w^\alpha,$$

where $\overset{\circ}{\Gamma}_{\mu\nu}^\alpha$ is the Riemannian connection defined in section 2 and $w^\alpha = g^{\alpha\mu} w_\mu$. Hence,

$$\gamma_{\mu\nu}^\alpha = \delta_\mu^\alpha w_\nu - g_{\mu\nu} w^\alpha.$$

Proposition 5.2. *The curvature tensors $\tilde{R}_{\mu\nu\sigma}^\alpha$, $\hat{R}_{\mu\nu\sigma}^\alpha$ and $\overset{\circ}{R}_{\mu\nu\sigma}^\alpha$ of the connections $\tilde{\Gamma}_{\mu\nu}^\alpha$, $\hat{\Gamma}_{\mu\nu}^\alpha$ and $\overset{\circ}{\Gamma}_{\mu\nu}^\alpha$ are expressed in terms of w_μ in the form:*

- (a) $\tilde{R}_{\mu\nu\sigma}^\alpha = \delta_\sigma^\alpha w_{\nu|\mu} - \delta_\nu^\alpha w_{\sigma|\mu}.$
- (b) $\hat{R}_{\mu\nu\sigma}^\alpha = \frac{1}{2}(\delta_\sigma^\alpha w_{\nu|\mu} - \delta_\nu^\alpha w_{\sigma|\mu}) + \frac{1}{4}w_\mu(\delta_\sigma^\alpha w_\nu - \delta_\nu^\alpha w_\sigma).$
- (c) $\overset{\circ}{R}_{\mu\nu\sigma}^\alpha = \delta_\mu^\alpha(w_{\nu|\sigma} - w_{\sigma|\nu}) + (g_{\mu\sigma}w_{\alpha|\nu} - g_{\mu\nu}w_{\alpha|\sigma}) + 2w^\alpha(g_{\mu\sigma}w_\nu - g_{\mu\nu}w_\sigma).$

Consequently, if $\tilde{R}_{\alpha\mu\nu\sigma} := g_{\alpha\epsilon}\tilde{R}_{\mu\nu\sigma}^\epsilon$ with similar expressions for $\hat{R}_{\alpha\mu\nu\sigma}$ and $\overset{\circ}{R}_{\alpha\mu\nu\sigma}$, we get:

- (á) $\tilde{R}_{\alpha\mu\nu\sigma} = g_{\alpha\sigma}w_{\nu|\mu} - g_{\alpha\nu}w_{\sigma|\mu}.$
- (b') $\hat{R}_{\alpha\mu\nu\sigma} = \frac{1}{2}(g_{\alpha\sigma}w_{\nu|\mu} - g_{\alpha\nu}w_{\sigma|\mu}) + \frac{1}{4}w_\mu(g_{\alpha\sigma}w_\nu - g_{\alpha\nu}w_\sigma).$
- (c') $\overset{\circ}{R}_{\alpha\mu\nu\sigma} = g_{\alpha\mu}(w_{\nu|\sigma} - w_{\sigma|\nu}) + (g_{\mu\sigma}w_{\alpha|\nu} - g_{\mu\nu}w_{\alpha|\sigma}) + 2w_\alpha(g_{\mu\sigma}w_\nu - g_{\mu\nu}w_\sigma).$

Proof. The first two relations hold by substituting $\Lambda_{\mu\nu}^\alpha$ in the formulae expressing $\tilde{R}_{\mu\nu\sigma}^\alpha$ (Theorem 3.1 (a)) and $\hat{R}_{\mu\nu\sigma}^\alpha$ (Proposition 3.8) respectively. The third relation holds by substituting $\Lambda_{\mu\nu}^\alpha$ and $\gamma_{\mu\nu}^\alpha$ in the formula expressing $\overset{\circ}{R}_{\mu\nu\sigma}^\alpha$ (Theorem 3.1 (c)). The remaining relations are straightforward. $\square$

Proposition 5.3. Let $\tilde{R}_{\mu\nu} := \tilde{R}_{\mu\nu\alpha}^\alpha$ and $\tilde{R} := g^{\mu\nu}\tilde{R}_{\mu\nu}$, with similar expressions for $\hat{R}_{\mu\nu\sigma}^\alpha$ and $\hat{R}_{\mu\nu\alpha}^\alpha$. Then

- (a) $\tilde{R}_{\mu\nu} = (n-1)w_{\nu|\mu}$, $\tilde{R} = (n-1)w^\mu{}_{|\mu}$.
- (b) $\hat{R}_{\mu\nu} = \frac{1}{4}(n-1)(2w_{\nu|\mu} + w_\mu w_\nu)$, $\hat{R} = \frac{1}{4}(n-1)(2w^\mu{}_{|\mu} + w^\mu w_\mu)$.
- (c) $\hat{R}_{\mu\nu}^\alpha = w_{\nu|\mu} - g_{\mu\nu}w^\sigma{}_{|\sigma} + 2(w_\nu w_\mu - g_{\mu\nu}w^\sigma w_\sigma)$, $\hat{R}^\alpha = (1-n)(w^\mu{}_{|\mu} + 2w^\mu w_\mu)$.

Proof. Follows directly from the relations obtained in Proposition 5.2 by applying the suitable contractions. $\square$

Theorem 5.4. The second order covariant tensor $w_{\nu|\mu}$ is symmetric:

$$w_{\mu|\nu} = w_{\nu|\mu}.$$

Proof. Follows directly from Proposition 5.3 (c) noting that $\hat{R}_{\mu\nu}$ is symmetric, being the Ricci tensor of the Riemannian connection. This can be also deduced from Theorem 5.2 (c) noting that $\hat{R}_{\mu\nu\sigma}^\mu = 0$. $\square$

A direct consequence of the above theorem is the following

Theorem 5.5. For an AP-space (M, λ) with semi-symmetric connection $\Gamma_{\mu\nu}^\alpha$, the following properties hold:

- (a) $\tilde{R}_{\mu\nu}$ and $\hat{R}_{\mu\nu}$ are symmetric.
- (b) $\tilde{R}_{\mu\nu\sigma}^\mu = 0$.
- (c) $\hat{R}_{\mu\nu\sigma}^\mu = 0$.

Theorem 5.6. For an AP-space (M, λ) with semi-symmetric connection $\Gamma_{\mu\nu}^\alpha$, the W-tensor has the form:

- (a) $W_{\mu\nu\sigma}^\alpha = \delta_\sigma^\alpha(w_{\nu|\mu} - w_\mu w_\nu) - \delta_\nu^\alpha(w_{\sigma|\mu} - w_\sigma w_\mu)$.
- (b) $W_{\alpha\mu\nu\sigma} = g_{\alpha\sigma}(w_{\nu|\mu} - w_\mu w_\nu) - g_{\alpha\nu}(w_{\sigma|\mu} - w_\sigma w_\mu)$ ($W_{\alpha\mu\nu\sigma} = g_{\alpha\epsilon}W_{\mu\nu\sigma}^\epsilon$).

Proof. Follows directly from Theorem 4.2 by substituting the expression of $\Lambda_{\mu\nu}^\alpha$ in terms w_μ . $\square$

Corollary 5.7. Let $W_{\mu\nu} := W_{\mu\nu\epsilon}^\epsilon$ and $W := g^{\mu\nu}W_{\mu\nu}$. Then

- (a) $W_{\mu\nu} = (n-1)(w_{\nu|\mu} - w_\mu w_\nu)$.
- (b) $W = (n-1)(w^\mu{}_{|\mu} - w^\mu w_\mu)$.
- (c) $W_{\mu\nu\sigma}^\mu = 0$.

Consequently, $W_{\mu\nu}$ is symmetric.

Comparing Corollary 5.7 (a) and Theorem 5.6 (a), we obtain

Theorem 5.8. *Let $\dim M \geq 2$. A sufficient condition for the vanishing of the W-tensor $W_{\mu\nu\sigma}^\alpha$ is that $W_{\mu\nu} = 0$.*

Finally, in view of Theorem 1 of K. Yano [18], the following result follows:

Theorem 5.9. *If the canonical connection $\Gamma_{\mu\nu}^\alpha$ of an AP-space is semi-symmetric, then the associated Riemannian metric $g_{\mu\nu}$ is conformally flat.*

Special case. In the following we assume that the canonical connection $\Gamma_{\mu\nu}^\alpha$ is a semi-symmetric connection whose defining 1-form w_μ satisfies the condition

$$w_\mu w^\nu = \delta_\mu^\nu.$$

It is easy to see that the above condition implies that $w_{\mu|\nu} = 0$ and that $w_\mu w_\nu = g_{\mu\nu}$. Under the given assumption, the different curvature tensors and the Wanas tensor, according to Proposition 5.2 and Theorems 5.6, take the form:

$$\tilde{R}_{\mu\nu\sigma}^\alpha = 0,$$

$$\hat{R}_{\mu\nu\sigma}^\alpha = \frac{1}{4}(\delta_\sigma^\alpha g_{\mu\nu} - \delta_\nu^\alpha g_{\mu\sigma}),$$

$$\mathring{R}_{\mu\nu\sigma}^\alpha = 2(\delta_\nu^\alpha g_{\mu\sigma} - \delta_\sigma^\alpha g_{\mu\nu}),$$

$$W_{\mu\nu\sigma}^\alpha = \delta_\nu^\alpha g_{\mu\sigma} - \delta_\sigma^\alpha g_{\mu\nu}.$$

Consequently,

$$W_{\mu\nu\sigma}^\alpha = -4\hat{R}_{\mu\nu\sigma}^\alpha = \frac{1}{2}\mathring{R}_{\mu\nu\sigma}^\alpha$$

and

$$\mathring{R}_{\alpha\mu\nu\sigma} = 2(g_{\alpha\nu}g_{\mu\sigma} - g_{\alpha\sigma}g_{\mu\nu}).$$

In this case, the W-tensor becomes a curvature-like tensor [8] and the above formulae imply the following result:

Theorem 5.10. *Let the canonical connection $\Gamma_{\mu\nu}^\alpha$ of an AP-space (M, λ) be a semi-symmetric connection whose defining 1-form w_μ satisfies the condition*

$$w_\mu w^\nu = \delta_\mu^\nu.$$

Then, all nonvanishing curvature tensors of (M, λ) coincide, up to a constant, with the W-tensor and the AP-space becomes a Riemannian space of constant curvature.

It should be noted that, in this particular case, the AP-character of the space recedes, whereas the Riemannian aspects of the AP-space become dominant. Physically speaking, the latter result seems to suggest that, in this particular case, electromagnetic effects are absent. This particular case can thus be considered, in some sense, as a limiting case.

References

- [1] O. C. Andonie et D. Smaranda, *Certaines connexions semi-symétrique*, Tensor, N.S., Vol. 31 (1977), 8-12.
- [2] F. Brickell and R. S. Clark, *Differentiable manifolds*, Van Nostrand Reinhold Co., 1970.
- [3] A. Einstein, *Unified field theory based on Riemannian metrics and distant parallelism*, Math. Annal., 102 (1930), 685-697.
- [4] A. Einstein, *The meaning of relativity*, 5th. Ed., Princeton Univ. Press, 1955.
- [5] K. Hayashi and T. Shirafuji, *New general relativity*, Phys. Rev. D 19 (1979), 3524-3554.
- [6] T. Imai, *Notes on semi-symmetric metric connections*, Tensor, N.S., Vol. 24 (1972), 293-296.
- [7] T. Imai, *Notes on semi-symmetric metric connections*, Tensor, N.S., Vol. 27 (1973), 56-58.
- [8] S. Kobayashi and K. Nomizu, *Foundations of differential geometry*, Vol. I. Interscience Publishers, 1963.
- [9] F. I. Mikhail and M.I. Wanas, *A generalized field theory, I. Field equations*, Proc. R. Soc. London, A. 356 (1977), 471-481.
- [10] F. I. Mikhail and M.I. Wanas, *A generalized field theory, II. Linearized field equations*, Int. J. Theoret. Phys., 20 (1981), 671-680.
- [11] H. P. Robertson, *Groups of motion in spaces admitting absolute parallelism*, Ann. Math, Princeton (2), 33 (1932), 496 - 520.
- [12] L. Tamássy, *Area and metrical connections in Finsler spaces, Finsler geometries*, A meeting of Minds, Kluwer Publ., 2000, Proc. of Coll. on Finsler Spaces, Edmonton, July 1998, 263-281.
- [13] L. Tamássy, *Geometry of the point Finsler spaces*, From the Book [Proceedings of J. Boliai Memorial Conference, Budapest, July 2006], "Non-Euclidean geometries", Springer, 2006, 445-461.
- [14] M. I. Wanas, *A generalized field theory and its applications in cosmology*, Ph.D Thesis, Cairo University, 1975.
- [15] M. I. Wanas, *A generalized field theory: Charged spherical symmetric solutions*, Int. J. Theoret. Phys., 24 (1985), 639-651.
- [16] M. I. Wanas, *A self consistent world model*, Astrophys. Space Sci., 154 (1989), 165-177.

- [17] M. I. Wanas, *Absolute parallelism geometry: Developments, applications and problems*, Stud. Cercet, Stiin. Ser. Mat. Univ. Bacau, No. 10 (2001), 297-309.
- [18] K. Yano, *On semi-symmetric metric connection*, Rev. Roumaine Math. Pures Appl., T. 15 (1970), 1579-1586.
- [19] N. L. Youssef, *Connexions métrique semi-symétriques semi-basiques*, Tensor, N.S., Vol. 40 (1983), 242-248.
- [20] N. L. Youssef, *Vertical semi-symmetric metric connections*, Tensor, N.S., Vol. 49 (1990), 218-229.